

MINISTRY OF EDUCATION, SINGAPORE
in collaboration with
CAMBRIDGE ASSESSMENT INTERNATIONAL EDUCATION
General Certificate of Education Ordinary Level

CANDIDATE
NAME



CENTRE
NUMBER

S				
---	--	--	--	--

INDEX
NUMBER

--	--	--	--

ADDITIONAL MATHEMATICS

4049/01

Paper 1

October/November 2023

2 hours 15 minutes

Candidates answer on the Question Paper.

No Additional Materials are required.

READ THESE INSTRUCTIONS FIRST

Write your Centre number, index number and name in the spaces at the top of this page.

Write in dark blue or black pen.

You may use an HB pencil for any diagrams or graphs.

Do not use staples, paper clips, glue or correction fluid.

DO NOT WRITE ON ANY BARCODES.

Answer **all** the questions.

Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place in the case of angles in degrees, unless a different level of accuracy is specified in the question.

The use of an approved scientific calculator is expected, where appropriate.

You are reminded of the need for clear presentation in your answers.

The number of marks is given in brackets [] at the end of each question or part question.

The total number of marks for this paper is 90.

This document consists of **19** printed pages and **1** blank page.



Singapore Examinations and Assessment Board



Cambridge Assessment
International Education



Mathematical Formulae

1. ALGEBRA

Quadratic Equation

For the equation $ax^2 + bx + c = 0$,

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

Binomial expansion

$$(a+b)^n = a^n + \binom{n}{1}a^{n-1}b + \binom{n}{2}a^{n-2}b^2 + \dots + \binom{n}{r}a^{n-r}b^r + \dots + b^n,$$

where n is a positive integer and $\binom{n}{r} = \frac{n!}{r!(n-r)!} = \frac{n(n-1)\dots(n-r+1)}{r!}$

2. TRIGONOMETRY

Identities

$$\sin^2 A + \cos^2 A = 1$$

$$\sec^2 A = 1 + \tan^2 A$$

$$\operatorname{cosec}^2 A = 1 + \cot^2 A$$

$$\sin(A \pm B) = \sin A \cos B \pm \cos A \sin B$$

$$\cos(A \pm B) = \cos A \cos B \mp \sin A \sin B$$

$$\tan(A \pm B) = \frac{\tan A \pm \tan B}{1 \mp \tan A \tan B}$$

$$\sin 2A = 2 \sin A \cos A$$

$$\cos 2A = \cos^2 A - \sin^2 A = 2 \cos^2 A - 1 = 1 - 2 \sin^2 A$$

$$\tan 2A = \frac{2 \tan A}{1 - \tan^2 A}$$

Formulae for ΔABC

$$\frac{a}{\sin A} = \frac{b}{\sin B} = \frac{c}{\sin C}$$

$$a^2 = b^2 + c^2 - 2bc \cos A$$

$$\Delta = \frac{1}{2}bc \sin A$$



- 1 Find the value of the constant c such that the line $y = 2x + c$ is a tangent to the curve $y = x^2 + 3x + 1$. [3]

Step 1: Let $x^2 + 3x + 1 = 2x + c$
 $x^2 + x + (1 - c) = 0$

Step 2: $b^2 - 4ac = 0$
 $1 - 4(1)(1 - c) = 0$
 $1 - 4 + 4c = 0$
 $4c = 3$
 $c = \frac{3}{4}$

- 2 Express $\frac{18 + 11x - 2x^2}{(x - 1)(x + 2)^2}$ in partial fractions. [5]

$$\frac{18 + 11x - 2x^2}{(x - 1)(x + 2)^2} = \frac{A}{(x - 1)} + \frac{B}{(x + 2)} + \frac{C}{(x + 2)^2}$$

By Cover Up Rule: Let $x = 1$

$$A = \frac{18 + 11 - 2}{3^2} = 3$$

$$\frac{18 + 11x - 2x^2}{(x - 1)(x + 2)^2} = \frac{3}{(x - 1)} + \frac{B}{(x + 2)} + \frac{C}{(x + 2)^2}$$

$$= \frac{3(x + 2)^2 + B(x - 1)(x + 2) + C(x - 1)}{(x - 1)(x + 2)^2}$$

By Substitution:

Let $x = -2$: $18 - 22 - 8 = C(-3)$
 $-12 = -3C$
 $\therefore C = 4$

Let $x = 0$: $18 = 3(4) + B(-1)(2) + 4(-1)$
 $18 = 12 - 2B - 4$
 $2B = -10$
 $B = -5$

$$\therefore \frac{18 + 11x - 2x^2}{(x - 1)(x + 2)^2} = \frac{3}{(x - 1)} - \frac{5}{(x + 2)} + \frac{4}{(x + 2)^2}$$



3 Prove that $\frac{\cos^2 \theta}{(\operatorname{cosec} \theta - 1)(\operatorname{cosec} \theta + 1)} + \frac{\sin^2 \theta}{(\sec \theta - 1)(\sec \theta + 1)} = 1.$ [5]

$$\begin{aligned}
 \text{LHS} &= \frac{\cos^2 \theta}{(\operatorname{cosec} \theta - 1)(\operatorname{cosec} \theta + 1)} + \frac{\sin^2 \theta}{(\sec \theta - 1)(\sec \theta + 1)} \\
 &= \frac{\cos^2 \theta}{\operatorname{cosec}^2 \theta - 1} + \frac{\sin^2 \theta}{\sec^2 \theta - 1} \\
 &= \frac{\cos^2 \theta}{\cot^2 \theta} + \frac{\sin^2 \theta}{\tan^2 \theta} \\
 &= \frac{\cos^2 \theta \tan^2 \theta + \cot^2 \theta \sin^2 \theta}{(\cot^2 \theta)(\tan^2 \theta)} \\
 &= \frac{\cancel{\cos^2 \theta} \left(\frac{\sin^2 \theta}{\cancel{\cos^2 \theta}} \right) + \left(\frac{\cancel{\cot^2 \theta}}{\cancel{\sin^2 \theta}} \right) \sin^2 \theta}{1} \\
 &= \sin^2 \theta + \cos^2 \theta \\
 &= 1 \\
 &= \text{RHS (proven)}
 \end{aligned}$$

Recap Identities:

- $1 + \tan^2 \theta = \sec^2 \theta$
- $1 + \cot^2 \theta = \operatorname{cosec}^2 \theta$

$$\begin{aligned}
 \cot \theta &= \frac{1}{\tan \theta} \\
 \therefore \cot^2 \theta &= \frac{1}{\tan^2 \theta}
 \end{aligned}$$



THE ANNEXE PROJECT
EDUCATIONAL CENTRE

ESTD 2008

The suggested solutions are prepared by Mr Alvin Yeo. Mr Yeo will hold no liability for any errors.





4 (a) Find $\frac{d}{dx}(xe^{-2x})$.

[3]

$$= x(-2e^{-2x}) + e^{-2x}(1)$$

$$= \underline{-2xe^{-2x} + e^{-2x}}$$

(b) Hence find $\int xe^{-2x} dx$.

[4]

from (a): $\int -2xe^{-2x} + e^{-2x} dx = xe^{-2x} + C$

$$-2 \int xe^{-2x} dx + \frac{e^{-2x}}{-2} = xe^{-2x} + C$$

$$\therefore -2 \int xe^{-2x} dx = \frac{1}{2}e^{-2x} + xe^{-2x} + C$$

$$\int xe^{-2x} dx = \frac{1}{4}e^{-2x} - \frac{1}{2}xe^{-2x} - \frac{1}{2}C$$

$$= \underline{\frac{1}{4}e^{-2x}(1+2x) + B}$$

(where $B = -\frac{1}{2}C$)



THE ANNEXE PROJECT
EDUCATIONAL CENTRE

ESTD 2008

The suggested solutions are prepared by Mr Alvin Yeo. Mr Yeo will hold no liability for any errors.





5 Solve the equation $\frac{\cos \theta + 4 \sin \theta}{2 \cos \theta + \sin \theta} = \cot \theta$ for $-\frac{\pi}{2} \leq \theta \leq \frac{\pi}{2}$.

[6]

$$\frac{\cos \theta + 4 \sin \theta}{2 \cos \theta + \sin \theta} = \frac{\cos \theta}{\sin \theta}$$

$$\cancel{\sin \theta \cos \theta} + 4 \sin^2 \theta = 2 \cos^2 \theta + \cancel{\sin \theta \cos \theta}$$

$$4 \sin^2 \theta - 2 \cos^2 \theta = 0$$

$$2 \sin^2 \theta - \cos^2 \theta = 0$$

$$2 \sin^2 \theta - (1 - \sin^2 \theta) = 0$$

$$3 \sin^2 \theta = 1$$

$$\sin^2 \theta = \frac{1}{3}$$

$$\sin \theta = \pm \frac{1}{\sqrt{3}}$$

Step 1: Basic angle for $\theta = 0.61548$ rad.

Step 2: $-\frac{\pi}{2} \leq \theta \leq \frac{\pi}{2}$

$$\therefore \theta = \underline{\underline{-0.615 \text{ or } 0.615}}$$



THE ANNEXE PROJECT
EDUCATIONAL CENTRE

ESTD 2008

The suggested solutions are prepared by Mr Alvin Yeo. Mr Yeo will hold no liability for any errors.



6 The function f is given by $f(x) = \frac{ax^2}{x-a}$, for $x > a$, where a is a positive constant.

(a) Find $f'(x)$.

[2]

$$\begin{aligned} f'(x) &= \frac{(x-a)(2ax) - (ax^2)(1)}{(x-a)^2} \\ &= \frac{2ax^2 - 2a^2x - ax^2}{(x-a)^2} \\ &= \frac{ax^2 - 2a^2x}{(x-a)^2} \\ &= \frac{ax(x-2a)}{(x-a)^2} \end{aligned}$$

The function g , defined for $x > a$, has the property that $g'(x) = (x-a)^2 f'(x)$.
 g decreases for $a < x < 8$.


(b) Find the value of a .

[4]

Step 1:

$$\begin{aligned} g'(x) &= \cancel{(x-a)^2} \cdot \frac{ax(x-2a)}{\cancel{(x-a)^2}} \\ &= ax(x-2a) \end{aligned}$$

Step 2: It is given that $g'(x) < 0$ for $a < x < 8$.
 and a is a positive constant



$$\begin{aligned} \text{Let } ax(x-2a) &< 0 \\ \therefore x(x-2a) &< 0 \\ 0 < x &< 2a \end{aligned}$$

Step 3: Since $x > a$ and $0 < x < 2a$,
 hence $a < x < 2a$.

$$\begin{aligned} \text{Comparing to } a < x < 8, \quad 2a &= 8 \\ \therefore a &= 4 \end{aligned}$$



- 7 Find the set of values of the constant k for which the curve $y = kx^2 + 4x + k - 3$ lies completely below the x -axis. [6]

Step 1: For a curve to lie completely below the x -axis, $k < 0$.

Step 2: When a curve lies completely below the x -axis, there are No Real Roots.

$$b^2 - 4ac < 0$$

$$4^2 - 4k(k-3) < 0$$


$$16 - 4k^2 + 12k < 0$$

$$\therefore k^2 - 3k - 4 > 0$$

$$(k-4)(k+1) > 0$$

(Divide both sides by -4)

Step 3:



From the graph, $k < -1$ or $k > 4$.

However, from step 1, $k < 0$.

Hence, $k < -1$



THE ANNEXE PROJECT
EDUCATIONAL CENTRE

ESTD 2008

The suggested solutions are prepared by Mr Alvin Yeo. Mr Yeo will hold no liability for any errors.





- 8 The line $y - 2x = 12$ intersects the curve $x^2 - xy + y^2 = 63$ at two points. Find the coordinates of these two points.

[5]

$$y = 2x + 12 \quad \text{--- (1)}$$

$$x^2 - xy + y^2 = 63 \quad \text{--- (2)}$$

Sub (1) into (2) :

$$x^2 - x(2x + 12) + (2x + 12)^2 = 63$$

$$x^2 - 2x^2 - 12x + 4x^2 + 48x + 144 - 63 = 0$$

$$3x^2 + 36x + 81 = 0$$

$$x^2 + 12x + 27 = 0$$

$$(x + 9)(x + 3) = 0$$

$$x = -3 \quad \text{or} \quad x = -9$$

$$\text{When } x = -3, \quad y = 2(-3) + 12 \\ = 6$$

$$\text{When } x = -9, \quad y = 2(-9) + 12 \\ = -6$$

The coordinates of the intersection points are $(-3, 6)$ and $(-9, -6)$



THE ANNEXE PROJECT
EDUCATIONAL CENTRE

ESTD 2008

The suggested solutions are prepared by Mr Alvin Yeo. Mr Yeo will hold no liability for any errors.





9 A curve has equation $y = 1 + x - \frac{x^2}{2} - 2x^3$.

(a) Find the x -coordinates of the stationary points, A and B , on the curve.

[4]

$$y = 1 + x - \frac{1}{2}x^2 - 2x^3$$

$$\frac{dy}{dx} = 1 - x - 6x^2$$

$$\text{Let } \frac{dy}{dx} = 0 : \quad -6x^2 - x + 1 = 0$$

$$6x^2 + x - 1 = 0$$

$$(3x-1)(2x+1) = 0$$

$$\therefore \underline{x = -\frac{1}{2} \text{ or } x = \frac{1}{3}}$$



THE ANNEXE PROJECT
EDUCATIONAL CENTRE

ESTD 2008

The suggested solutions are prepared by Mr Alvin Yeo. Mr Yeo will hold no liability for any errors.



DO NOT WRITE IN THIS MARGIN

It is given that P is a point on the curve where the gradient is a maximum.

(b) Show that P and the midpoint of AB have the same x -coordinate.

[4]

Let gradient be G :

Step 1: $G = 1 - x - 6x^2$

$$\frac{dG}{dx} = -1 - 12x$$

Step 2: Let $\frac{dG}{dx} = 0$

$$\therefore -1 - 12x = 0$$

$$12x = -1$$

$$x = -\frac{1}{12}$$

Step 3: $\frac{d^2G}{dx^2} = -12 < 0$

2nd Derivative Test

so when $x = -\frac{1}{12}$, gradient is maximum.

i.e. x -coordinate of $P = -\frac{1}{12}$

$$\begin{aligned} x\text{-coordinate of midpoint of } AB &= \frac{-\frac{1}{2} + \frac{1}{3}}{2} \\ &= -\frac{1}{12} \text{ (shown)} \end{aligned}$$



THE ANNEXE PROJECT
EDUCATIONAL CENTRE

ESTD 2008

The suggested solutions are prepared by Mr Alvin Yeo. Mr Yeo will hold no liability for any errors.





10 A circle, with centre C , has equation $x^2 + y^2 + 10x - 24y = 0$.

(a) Find the coordinates of C and the radius of the circle.

[4]

$$\begin{aligned}x^2 + 10x + y^2 - 24y &= 0 \\(x + 5)^2 - 5^2 + (y - 12)^2 - 12^2 &= 0 \\(x + 5)^2 + (y - 12)^2 &= 13^2\end{aligned}$$

$$\therefore C = (-5, 12) \text{ and radius } r = 13$$

(b) Find the coordinates of the points at which the circle intersects the y -axis.

[2]

$$\begin{aligned}\text{Let } x = 0: \quad y^2 - 24y &= 0 \\y(y - 24) &= 0 \\y = 0 \text{ or } y = 24\end{aligned}$$

$$\therefore \text{coordinates are } \underline{(0, 0) \text{ and } (0, 24)}.$$

The suggested solutions are prepared by Mr Alvin Yeo. Mr Yeo will hold no liability for any errors.




The point X is on the line which passes through C and the origin, O . It is given that the distance CX is three times the distance OC .

- (c) Find the coordinates of the possible positions of X .

[4]

Using Ratio Theorem:

$X_1(a, b)$




$$\frac{3(0) + 1(a)}{4} = -5 \quad \frac{3(0) + 1(b)}{4} = 12$$


$$\therefore a = -20$$

$$\therefore b = 48$$

Hence, $X_1 = (-20, 48)$



$C(-5, 12)$



$$\frac{1(c) + 2(-5)}{3} = 0$$

$$\frac{1(d) + 2(12)}{3} = 0$$

$$\therefore c = 10$$

$$\therefore d = -24$$

Hence, $X_2 = (10, -24)$

The suggested solutions are prepared by Mr Alvin Yeo. Mr Yeo will hold no liability for any errors.





- 11 A circular patch of water of negligible thickness is expanding on a thin piece of paper. At time t seconds the rate of change of the radius, r cm, of the patch is given by

$$\frac{dr}{dt} = \frac{k}{2t+1} \text{ cm/s, where } k \text{ is a constant.}$$

Initially the radius of the patch is 1 cm and the radius is increasing at a rate of 0.5 cm/s.

- (a) Show that $k = 0.5$.

$$\text{Let } 0.5 = \frac{k}{2(0)+1}$$

$$\therefore \underline{k = 0.5}$$

[1]

- (b) Find an expression for r in terms of t .

$$\begin{aligned} r &= \int \frac{0.5}{2t+1} dt \\ &= 0.5 \int \frac{1}{2t+1} dt \\ &= 0.5 \left[\frac{\ln(2t+1)}{2} \right] + C \\ &= \frac{1}{4} \ln(2t+1) + C \end{aligned}$$

[4]

Since $r = 1$ cm when $t = 0$,

$$\therefore 1 = \frac{1}{4} \ln 1 + C$$

$$C = 1$$

$$\text{Hence, } \underline{r = \frac{1}{4} \ln(2t+1) + 1}$$



THE ANNEXE PROJECT
EDUCATIONAL CENTRE

ESTD 2008

The suggested solutions are prepared by Mr Alvin Yeo. Mr Yeo will hold no liability for any errors.



- (c) Hence find the rate of increase of the area of the patch after the patch has been expanding for 3 seconds. [4]

Step 1: $A = \pi r^2$
 $\frac{dA}{dr} = 2\pi r$

Step 2: When $t = 3\text{ s}$, $r = \frac{1}{2} \ln(6+1) + 1$
 $= 1.4865\text{ cm.}$

Step 3: $\frac{dA}{dt} = \frac{dA}{dr} \times \frac{dr}{dt}$
 $= 2\pi(1.4865) \times \frac{0.5}{2(3)+1}$
 $= 0.66713$
 $= \underline{0.667\text{ cm}^2/\text{s}}$



THE ANNEXE PROJECT
 EDUCATIONAL CENTRE

ESTD 2008

The suggested solutions are prepared by Mr Alvin Yeo. Mr Yeo will hold no liability for any errors.





12 A ball is thrown vertically upwards. Its height, h m, above the ground at time t seconds after being thrown is given by the formula $h = 1.75 + 5t - 5t^2$.

(a) State the height above the ground from which the ball is thrown. [1]

$$\text{When } t = 0 \text{ s, } \underline{h = 1.75 \text{ m}}$$

(b) Express h in the form $a + b(t+c)^2$ where a , b and c are constants to be determined. [3]

$$\begin{aligned} h &= -5(t^2 - t) + 1.75 \\ &= -5\left[\left(t - \frac{1}{2}\right)^2 - \left(\frac{1}{2}\right)^2\right] + 1.75 \\ &= -5\left(t - \frac{1}{2}\right)^2 + 1.25 + 1.75 \\ &= \underline{3 - 5\left(t - \frac{1}{2}\right)^2} \end{aligned}$$

$$\text{where } a = 3, b = -5 \text{ and } c = -0.5$$

(c) Hence state the maximum height attained by the ball and the time at which this occurs. [2]

$$\begin{aligned} \text{Max. height} &= \underline{3 \text{ m}} \\ \text{when } t &= \underline{0.5 \text{ s}} \end{aligned}$$

The suggested solutions are prepared by Mr Alvin Yeo. Mr Yeo will hold no liability for any errors.



- (d) The ball hits the ground. Explain why the time taken for the ball to hit the ground is **not** twice the time found in part (c). [1]

Because the ball wasn't thrown up from the ground level. It was thrown from an initial height of 1.75m.

- (e) Find the length of time for which the ball is at least 2m above the ground. [3]

Step 1: Let $h = 2$

$$\therefore 3 - 5\left(t - \frac{1}{2}\right)^2 = 2$$

$$5\left(t - \frac{1}{2}\right)^2 = 1$$

$$t - \frac{1}{2} = \pm \sqrt{\frac{1}{5}}$$

$$t = \frac{1}{2} \pm \sqrt{\frac{1}{5}}$$

$$= \underline{0.052786 \text{ s} \text{ or } 0.94721 \text{ s}}$$

Step 2:

$$\text{duration} = 0.94721 - 0.052786$$

$$= \underline{0.894 \text{ s}}$$



THE ANNEXE PROJECT
EDUCATIONAL CENTRE

ESTD 2008

The suggested solutions are prepared by Mr Alvin Yeo. Mr Yeo will hold no liability for any errors.






13 The perpendicular bisector of the line joining the points $A(2, h)$ and $B(-8, -7)$ passes through the point $X(h, -\frac{13}{2})$, where h is a constant.

(a) Express the gradient of the perpendicular bisector of the line AB in terms of h . [3]

$$\begin{aligned} \text{Step 1: gradient } AB &= \frac{h - (-7)}{2 - (-8)} \\ &= \frac{h + 7}{10} \end{aligned}$$



$$\text{Step 2: gradient of perp. bisector} = \frac{-10}{h + 7}$$

for perpendicular lines,
 $m_1 \times m_2 = -1$

(b) Hence find the possible values of h . [7]

$$\begin{aligned} \text{Step 1:} \\ \text{midpoint of } AB &= \left(\frac{-8+2}{2}, \frac{-7+h}{2} \right) \\ M &= \left(-3, \frac{h-7}{2} \right) \end{aligned}$$

$$\begin{aligned} \text{Step 2:} \\ \text{gradient } MX &= \frac{-\frac{13}{2} - \left(\frac{h-7}{2}\right)}{h - (-3)} = \frac{-13 - (h-7)}{2(h+3)} \\ &= \frac{-6-h}{2h+6} \end{aligned}$$



THE ANNEXE PROJECT
EDUCATIONAL CENTRE

ESTD 2008

The suggested solutions are prepared by Mr Alvin Yeo. Mr Yeo will hold no liability for any errors.





Continuation of working space for question 13(b).

Step 3 :

$$\frac{-6-h}{2h+6} = \frac{-10}{h+7}$$

$$(-6-h)(h+7) = -10(2h+6)$$

$$-6h - 42 - h^2 - 7h = -20h - 60$$

$$h^2 - 7h - 18 = 0$$

$$(h+2)(h-9) = 0$$

$$\therefore \underline{h = -2 \text{ or } 9}$$



THE ANNEXE PROJECT
EDUCATIONAL CENTRE

ESTD 2008

The suggested solutions are prepared by Mr Alvin Yeo. Mr Yeo will hold no liability for any errors.







BLANK PAGE



DO NOT WRITE IN THIS MARGIN



Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

Cambridge Assessment International Education is part of Cambridge Assessment. Cambridge Assessment is the brand name of the University of Cambridge Local Examinations Syndicate (UCLES), which is a department of the University of Cambridge.

